

Algebraic And Partial Fractions

Algebraic Fractions

A polynomial is a sum of powers of a variable x , such as $x^3 - 3x^2 - 2x + 1$. An algebraic fraction is the ratio of two polynomials. If the numerator of a fraction is greater than the denominator, it is *improper*, otherwise it is *proper*.

Partial Fractions By Negating Variables

The expression $\frac{2}{x-1} + \frac{5}{x+4}$ can be combined into a single fraction by introducing the common denominator $(x-1)(x+4)$, which results in $\frac{7x+3}{(x-1)(x+4)}$. The method of partial fractions provides a method for reversing that operation.

1. Split the fraction based on the denominator, with A and B as the numerators.

$$\frac{7x+3}{(x-1)(x+4)} = \frac{A}{x-1} + \frac{B}{x+4} \quad (1)$$

2. Multiply the missing denominator portion.

$$= \frac{A(x+4)}{(x-1)(x+4)} + \frac{B(x-1)}{(x-1)(x+4)} \quad (2)$$

3. Simplify to a single equation.

$$= \frac{A(x+4) + B(x-1)}{(x-1)(x+4)} \quad (3)$$

4. Equate numerators.

$$7x+3 = A(x+4) + B(x-1) \quad (4)$$

5. Set one set of numerators equal to 0:

when $x = 1$:

6. And solve to get a value for one of the variables.

$$7 \cdot 1 + 3 = A(1+4) + B(1-1) \quad (5)$$

$$10 = 5A \therefore A = 2 \quad (6)$$

7. Set the other numerators equal to 0...

when $x = -4$:

8. And solve for the second value.

$$7 \cdot -4 + 3 = A(-4+4) + B(-4-1) \quad (7)$$

$$-25 = -5B \therefore B = 5 \quad (8)$$

9. Which gives a solution.

$$\frac{7x+3}{(x-1)(x+4)} = \frac{2}{x-1} + \frac{5}{x+4} \quad (9)$$

Examples

$$\begin{aligned}\frac{5+2}{(x+2)(5x-2)} &= \frac{A}{x+2} + \frac{B}{3x-2} \\ &= \frac{A(3x-2)}{(x+2)(3x-2)} + \frac{B(x+2)}{(x+2)(3x-2)} \\ 5x+2 &= A(3x-2) + B(x+2)\end{aligned}$$

when $x = -2$:

$$\begin{aligned}5 \cdot -2 + 2 &= A(3 \cdot -2 - 2) + B(-2 + 2) \\ -8 &= -8A \\ A &= 1\end{aligned}$$

when $x = \frac{2}{3}$:

$$\begin{aligned}5 \cdot \frac{2}{3} + 2 &= A(3 \cdot \frac{2}{3} - 2) + B(\frac{2}{3} + 2) \\ \frac{16}{3} &= \frac{8B}{3} \\ B &= 2\end{aligned}$$

$$\therefore \frac{5x+2}{(x+2)(3x-2)} = \frac{1}{x+2} + \frac{2}{3x-2}$$

$$\begin{aligned}\frac{3x+1}{(x-1)(x+1)} &= \frac{A}{x-1} + \frac{B}{x-1} \\ &= \frac{A(x+1)}{(x-1)(x+1)} + \frac{A(x-1)}{(x-1)(x+1)} \\ 3x+1 &= A(x+1) + B(x-1)\end{aligned}$$

when $x = 1$:

$$\begin{aligned}3 \cdot 1 + 1 &= A(1+1) + B(1-1) \\ 4 &= 2A \\ A &= 2\end{aligned}$$

when $x = -1$:

$$\begin{aligned}3 \cdot -1 + 1 &= A(-1+1) + B(-1-1) \\ -2 &= -2B \\ B &= 1\end{aligned}$$

$$\therefore \frac{3x+1}{(x-1)(x+1)} = \frac{2}{x-1} + \frac{1}{x+1}$$

Partial Fractions By Stratifying Exponents

Another method works by equating exponents.

Steps 1-4 are the same as above.	$7x + 3 = A(x + 4) + B(x - 1)$	
	$= Ax + 4A + Bx - B$	(10)
10. Expand both sides of the equation.	$= (A + B)x + (4A - B)$	(11)
11. Factor by powers of x.	$[x^1] : 7 = A + B$	(12)
12. Separate by powers of x to produce separate equations.	$[x^0] : 3 = 4A - B$	
	$7 + 3 = A + B + 4A - B$	(13)
13. Solve as simultaneous equations...	$10 = 5A$	
14. ...to find one of the two variables.	$A = 2$	(14)
15. Take one of the equations from Step 7, and solve for the other variable.	$7 = A + B$	(15)
	$= 2 + B$	
16. Which brings us to the same answer.	$B = 5$	
	$\therefore \frac{7x + 3}{(x - 1)(x + 4)} = \frac{2}{x - 1} + \frac{5}{x + 4}$	(16)

Examples

$$\begin{aligned} \frac{10x - 7}{(x - 4)(2x^2 - 6x + 3)} &= \frac{A}{x + 4} + \frac{Bx + C}{2x^2 - 6x} \\ &= \frac{A(2x^2 - 6x + 3)}{(x - 4)(2x^2 - 6x + 3)} + \frac{(Bx + C)(x + 4)}{(x - 4)(2x^2 - 6x + 4)} \\ 10x - 7 &= A(2x^2 - 6x + 3) + (Bx + C)(x + 4) \end{aligned}$$

when $x = 4$: $10 \cdot 4 - 7 = A(2 \cdot 4^2 - 6 \cdot 4 + 3) + (B \cdot 4 + C)(4 - 4)$
 $33 = A(32 - 24 + 3)$
 $A = 3$

$$\begin{aligned} 10x - 7 &= 2Ax^2 - 6Ax + 3A + Bx^2 - 4Bx + Cx - 4C \\ [x^2] : 0 &= 2A + B \\ [x^1] : 10 &= -6A - 4B + C \\ [x^0] : -7 &= 3A - 4C \\ 0 &= 2A + B \\ &= 2 \cdot 3 + B \\ B &= -6 \end{aligned}$$

$$\begin{aligned} 10 &= -6A - 4B + C \\ &= -6 \cdot 3 - 4 \cdot -6 + C \\ &= -18 + 10 - 24 \\ C &= 18 + 10 - 24 \\ &= 4 \end{aligned}$$

$$\frac{10x - 7}{(x - 4)(2x^2 - 6x + 3)} = \frac{3}{x - 4} + \frac{4 - 6x}{2x^2 - 6x + 3}$$

Combining Methods

It may be necessary to use both techniques to solve more complex fractions. This can be seen when solving the equation $\frac{9x+3}{(x-1)(x^2+x+10)} = \frac{A}{x-1} + \frac{Bx+C}{x^2+x+10}$.

$$9x + 3 = A(x^2 + x + 10) + (Bx + C)(x - 1) \quad (17)$$

when $x = 1$:

$$9 \cdot 1 + 3 = A(1^2 + 1 + 10) + B(\cdot 1 + C)(1 - 1)$$

$$12 = 12A$$

$$A = 1 \quad (18)$$

17. Use Steps 1-4 above.

18. Use Steps 5-6 to find A.

19. Use Steps 12-14 to find B.

20. Use Steps 5-6 again to find C.

$$9x + 3 = Ax^2 + Ax + 10A + Bx^2 - Bx + Cx - C$$

$$[x^2] : 0 = A + B$$

$$[x^1] : 9 = A - B + C$$

$$[x^0] : 3 = 10A - C$$

$$B = -A$$

$$= -1 \quad (19)$$

$$\text{when } x = 0 : 9 \cdot 0 + 3 = A(0^2 + 0 + 10) + (B \cdot 0 + C)(0 - 1)$$

$$3 = 10 \cdot A + C \cdot -1$$

$$3 = 10 - C$$

$$C = 7 \quad (20)$$

$$\therefore \frac{9x+3}{(x-1)(x^2+x+10)} = \frac{1}{x-1} + \frac{-x+7}{x^2+x+10} \quad (21)$$